

学校编码：10384

学号：2007171108

廈門大學

博士学位论文

微囊藻毒素 - LR在斑马鱼体内慢性毒性机制的研究

Toxicological study on the mechanism of
chronic microcystin-LR toxicity in
zebrafish (*Danio rerio*)

王明华

指导教师：王大志

专业名称：环境科学

答辩日期：2012年5月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1. 经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

摘 要

微囊藻毒素-LR (缩写为MCLR) 为水体中毒性最强、最经常爆发的一种毒素, 该毒素对生物体呈多种毒性效应 (如肝毒性和神经毒性), 然而其毒性作用机制至今还不清楚。鉴于此, 本研究工作以斑马鱼*D. rerio*为研究对象, 进行不同低浓度MCLR (空白对照组、2和20 $\mu\text{g/L}$ 毒素处理组, 暴露时间为30 d) 慢性胁迫下鱼体内肝脏和大脑的蛋白质组学研究; 即暴露30 d后, 取斑马鱼的肝脏和大脑, 采用双向凝胶电泳 (Two-dimensional electrophoresis, 简称为2-DE) 和基质辅助电离解析飞行时间质谱 (Matrix-assisted laser desorption/ionization time-of-flight/time-of-flight mass spectrometer, 简称为MALDI-TOF/TOF-MS) 等方法, 研究MCLR胁迫下两种组织蛋白质表达图谱所发生的变化, 找出发生显著变化的差异蛋白点, 对其进行功能鉴定, 继而揭示了由MCLR毒性引起的蛋白质组的变化; 另外, 本文测定了肝脏和大脑对MCLR的累积量及其蛋白磷酸酶 (Protein phosphatase, 简称为PP) 的活性, 同时也从超微结构水平来探讨MCLR所诱导的肝脏细胞损伤程度。该研究工作旨在探讨MCLR对鱼类的慢性毒害效应及其毒性机制, 同时为可特异地指示MCs污染的生物标志物的筛选提供理论和实践依据。主要研究结果如下:

- 1、暴露结束后, 毒素胁迫显著提高处理组斑马鱼肝脏内MCLR的累积量和PP活性, 如空白对照组中没有检测到MCLR的含量, 而2和20 $\mu\text{g/L}$ 毒素处理组的MCLR含量分别为0.031和0.039 $\mu\text{g/mg DW}$; 同样地, 相对于对照空白组, 20 $\mu\text{g/L}$ MCLR处理组的PP活性显著提高2倍。有趣的是, 免疫印记实验表明肝脏内蛋白磷酸酶2A (缩写为PP2A) 含量与MCLR处理浓度无相关性。MCLR胁迫引起肝脏细胞超微结构的显著损伤: 相对于空白对照组, 处理组肝脏细胞内粗内质网发生普遍的肿胀, 伴随着网槽的扩张或空泡化; 线粒体也发生肿胀, 许多线粒体失去嵴; 最为明显的是, 处理组肝脏细胞有蜂窝状结构体的出现。蛋白质组学分析结果表明有22个蛋白质斑点在毒素胁迫下发生显著的变化; 经过鉴定, 这些蛋白质参与细胞骨架的组装、大分子的代谢、氧化胁迫和信号传导; 因此, MCLR慢性肝毒性可引起细胞骨架组装和生物大分子代谢的失常, 诱导氧化胁迫的出现, 伴随着细胞内信

号传导的扰乱。另外，MCLR慢性肝毒性可能启动活性氧（缩写为ROS）途径，而不是PP途径，后者为该毒素急性毒害效应的主要机制。

2、暴露实验结束后，毒素胁迫显著提高斑马鱼大脑内MCLR含量和PP活性，如空白对照组中没有检测到MCLR的累积，而2和20 $\mu\text{g/L}$ MCLR处理组的毒素含量分别为0.030和0.053 $\mu\text{g/mg DW}$ ；同样地，相对于对照空白组，20 $\mu\text{g/L}$ MCLR处理组的PP活性显著提高1.3倍。蛋白质组学分析结果表明大脑内有30个蛋白质斑点在毒素胁迫下发生显著的变化；经过鉴定，这些蛋白质参与细胞骨架的组装、生物大分子的代谢、氧化胁迫、信号传导、以及其它功能（如转运、蛋白质降解、细胞凋亡和蛋白质翻译）。综上所述，MCLR的神经毒性可引起氧化胁迫、细胞骨架组装和大分子代谢的异常，以及伴随着信号传导和其它功能（如蛋白质的降解、细胞转运、细胞凋亡和蛋白质翻译）的紊乱，暗示该毒素对斑马鱼大脑神经毒性的复杂性和多样性。斑马鱼大脑的PP活性随着MCLR浓度的增加而升高，可能与毒素胁迫下PP2C β 的诱导表达相关。另外，该研究提示MCLR的慢性神经毒性启动于ROS和PP途径，然而PP途径与PP2C β 的显著上调相关的。最后，MCLR可能具有内分泌干扰物质的效应，因为该毒素胁迫显著地诱导卵黄蛋白原的形成；然而具体原因和机制的探讨有待于进一步研究。

值得一提的是，MCLR肝脏毒性和神经毒性引起的蛋白质组变化很相似，即两种毒性均显著影响了参与细胞代谢、骨架蛋白组装、氧化胁迫和信号传导的细胞过程，说明了MCLR的多种毒性效应在某种程度上呈一致性，虽然MCLR神经毒性还特异地影响了细胞转运、蛋白质降解、细胞凋亡和蛋白质翻译的生化过程；然而，即使MCLR两种毒性影响了相似的细胞过程（如细胞代谢、细胞骨架蛋白的组装、氧化胁迫和信号传导），具体参与的蛋白质却不相同，暗示该毒素肝毒性和神经毒性的毒害效应可能存在不同的生化机制。总之，本研究清楚地展现蛋白质组学技术可有效地对MCLR在斑马鱼体内肝毒性和神经毒性的毒害效应进行一些机制上的探讨。

关键词：蛋白磷酸酶；蛋白质组学；肝毒性；活性氧；MCLR；神经毒性

Abstract

Microcystin-LR (MCLR) is the most toxic and most frequently encountered toxin in the aquatic environment. MCLR presents multi-toxicity (e.g. hepatotoxicity and neurotoxicity), however the exact mechanism of which is still unknown. So, this study investigated the protein profiles of zebrafish (*Danio rerio*) liver or brain chronically exposed to MCLR concentrations (2 or 20 $\mu\text{g/L}$) using the proteomic approach. Namely, after 30 d MCLR exposure, under the cooperation of two-dimensional electrophoresis (2-DE) and matrix-assisted laser desorption/ionization time-of-flight/time-of-flight mass spectrometer (MALDI-TOF/TOF-MS) analysis, this work examined the differential protein profiles of both the tissues (liver or brain), revealed the differential protein spots, and subsequently submitted them to MS identity, with being aimed at elucidating the proteomic change caused by MCLR toxicity. Also, this work investigated toxin accumulation and protein phosphatase (PP) activity in both the tissues under different MCLR treatments. Meanwhile, this study especially analyzed the MCLR-induced damage in the hepatocytes at the ultrastructural level. Taken together, this work provides a new insight into MCLR toxicity and its toxic mechanism in fish, and helps to select out a potential biomarker for biomonitoring MCs pollution in the aquatic environment. The results of the present study are described as follows:

1. MCLR treatment significantly enhanced toxin accumulation and PP activity in the treated zebrafish livers at the end of this exposure experiment. For example, no MCLR was detected in MCLR non-exposed zebrafish livers, while the toxin contents attained to 0.031 and 0.039 $\mu\text{g/mg DW}$, respectively, in the zebrafish livers exposed to 2 and 20 $\mu\text{g/L}$ MCLR. Similarly, in comparison to the control, the hepatic PP activity increased two times in zebrafish livers exposed to 20 $\mu\text{g/L}$ MCLR. However, using the antibody representing

a specific interaction with PP2A, the Western blot analysis indicated that the hepatic PP2A amount was independent of the ambient MCLR concentrations. MCLR caused a noticeable damage to liver ultrastructure, a widespread swelling in the rough endoplasmatic reticulum and mitochondria was observed in the MCLR-exposed hepatocytes, and a honeycomb-like structure was formed in the treated nucleoli. The proteomic analysis revealed that the abundance of 22 protein spots, measured by 2-DE, was remarkably altered in response to toxin exposure. These proteins were involved in cytoskeleton assembly, macromolecule metabolism, oxidative stress and signal transduction. Thus, the chronic effects of MCLR hepatotoxicity caused dysfunction of cytoskeleton assembly and macromolecule metabolism, and induced oxidative stress with a concomitant interference with cell signal transduction. We speculate that the chronic effects of MCLR hepatotoxicity might initiate the ROS pathway, instead of the PP pathway which is the main mechanism for MCLR acute toxicity.

2. MCLR treatment obviously enhanced toxin accumulation and PP activity in the treated zebrafish brains after the 30 d exposure. For instance, no MCLR was detected in the control while the toxin contents were 0.030 and 0.053 $\mu\text{g}/\text{mg}$ DW in the brains exposed to 2 and 20 $\mu\text{g}/\text{L}$ MCLR. Similarly, compared to the control, the brain PP activity increased 1.3 times in zebrafish brains exposed to 20 $\mu\text{g}/\text{L}$ MCLR. The proteomic analysis indicated that the abundance of 30 protein spots was remarkably altered in response to MCLR exposure. These proteins are involved in cytoskeleton assembly, macromolecule metabolism, oxidative stress, signal transduction, and other functions (e.g. transporting, protein degradation, apoptosis and translation). Overall, MCLR neurotoxicity induced oxidative stress, and a dysfunction of cytoskeleton assembly and macromolecule metabolism, with a concomitant interference with signal transduction and other functions (e.g. protein degradation, transport, apoptosis and translation), suggesting that MCLR toxicity to the zebrafish brain

was complex and diverse. Also, the PP activity in the brain was enhanced with an increasing MCLR concentration, and this was partly exemplified by an overexpression of PP2C^{1537;2} under toxin treatment. Thus, our study firstly demonstrated that the chronic neurotoxicity of MCLR might initiate the PP pathway via an upregulation of PP2C in the zebrafish brain, in addition to the ROS pathway. Interestingly, an increased vitellogenin expression in the treated group suggested that MCLR might mimic the effects of endocrine disruption compounds, and this really deserves further study.

It should be noted that the responses in brain were quite similar with liver in our study, i.e. both toxicities exerted a significant effect on the similar cellular processes overlapped in general functional categories (e.g. metabolism, cytoskeletal assembly, oxidative stress and signal transduction), highlighting that to some extent, there is a similarity in MCLR multi-toxicities, although MCLR neurotoxicity specifically affected transport, protein degradation, apoptosis and translation in zebrafish brains. However, even though the affected cellular processes were overlapped in general functional categories (e.g. metabolism, cytoskeletal assembly, oxidative stress and signal transduction), none of the proteins in the functional category in brain was the same as that from liver., and a different biochemical mechanism might be involved in the initiation of these two toxicities. Taken together, proteomics provides a potential tool for studying the biochemical mechanisms concerning MCLR toxicity in zebrafish.

Keywords: Protein phosphatase; Proteomics; Hepatotoxicity; Reactive oxygen species; MCLR; Neurotoxicity

参考资料

- Allen DA, Ottaway JH, 1986. Succinate thiokinase in pigeon breast muscle mitochondria. *FEBS Lett.*, 194: 171 – 175.
- Ame MV, Baroni MV, Galanti LN, Bocco JL, Wunderlin DA, 2009. Effects of microcystin – LR on the expression of P – glycoprotein in *Jenynsia multidentata*. *Chemosphere*, 74: 1179 – 1186.
- Andersen L, Goto-Kazeto R, Trant JM, Nash JP, Korsgaard B, Bjerregaard P, 2006. Short-term exposure to low concentrations of the synthetic androgen methyltestosterone affects vitellogenin and steroid levels in adult male zebrafish (*Danio rerio*). *Aquat. Toxicol.*, 76: 343 – 352
- Ariza RR, Keyse SM, Moggs JG, Wood RD, 1996. Reversible protein phosphorylation modulates nucleotide excision repair of damaged DNA by human cell extracts. *Nucl. Acid. Res.*, 24: 433 – 440.
- Asanuma M, Ogawa N, Hirata H, Chou H, Kondo Y, Mori A, 1993. Ischemia-induced changes in β -tubulin and α -actin mRNA in the gerbil brain and effects of bifemelane hydrochloride. *Brain Res.*, 600: 243 – 248.
- Babior BM, 2000. Phagocytes and oxidative stress. *Am. J. Med.*, 109: 33 – 44.
- Bard SM, 2000. Multixenobiotic resistance as a cellular defense mechanism in aquatic organisms. *Aquat. Toxicol.*, 48: 357 – 389.
- Barker FG, Chen P, Furman F, Aldape KD, Edwards MS, Israel MA, 1997. P16 deletion and mutation analysis in human brain tumors. *J. Neurooncol.*, 31: 17 – 23.
- Bernardi P, Scorrano L, Colonna R, Petronilli V, Di Lisa F, 1999. Mitochondria and cell death. Mechanistic aspects and methodological issues. *Eur. J. Biochem.*, 264: 687 – 701.
- Betty M, Harnish SW, Rhodes KJ, Cockett MI, 1998. Distribution of heterotrimeric G – protein β and γ subunits in the rat brain. *Neuroscience*, 85: 475 – 486.
- Blom JF, Jüttner F, 2005. High crustacean toxicity of microcystin congeners does not correlate with high protein phosphatase inhibitory activity. *Toxicon*, 46: 465 – 470.
- Booth MPS, Connors R, Rumsby G, Brady RL, 2006. Structural basis of substrate specificity in human glyoxylate reductase/hydroxypyruvate reductase. *J. Mol. Biol.*, 360: 178 – 189.
- Boscoboinik D, Szewczyk A, Hensey C, Azzi A, 1991. Inhibition of cell proliferation by α -tocopherol. Role of protein kinase C. *J. Biol. Chem.*, 266: 6188 – 6194.
- Burke WJ, Li SW, Williams EA, Nonneman R, Zahm DS, 2003. 3,4-Dihydroxyphenylacetaldehyde is the toxic dopamine metabolite in vivo: implications for Parkinson ' s disease pathogenesis. *Brain Res.*, 989: 205 – 213.
- Burridge K, Feramisco JR, 1981. Non – muscle β – actinins are calcium-sensitive actin-binding proteins. *Nature*, 294: 565 – 567.
- Cabrera-Vera TM, Vanhauwe J, Thomas TO, Medkova M, Preininger A, Mazzoni MR, Hamm HE, 2003. Insights into G protein structure, function, and regulation. *Endocr. Rev.*, 24: 765 – 781.
- Campos A, Vasconcelos V, 2010. Molecular mechanisms of microcystin toxicity in animal cells. *Int. J. Mol. Sci.*, 11: 268 – 287.
- Carmichael WW, Azevedo SM, An JS, Molica RJ, Jochimsen EM, Lau S, Rinehart KL, Shaw GR, Eaglesham GK, 2001. Human fatalities from cyanobacteria: chemical and biological evidence for cyanotoxins. *Environ. Health Perspect.* 109: 663 – 668.
- Carlsson L, Nyström LE, Lindberg U, Kannan KK, Cid-Dresdner H, Lövgren S, Jönvall H, 1976. Crystallization of a non-muscle actin. *J. Mol. Biol.*, 105: 353 – 366.
- Cazenave J, Wunderlin DA, Bistoni MA, Amé MV, Krause E, Pflugmacher S, Wiegand C, 2005. Uptake, tissue distribution and accumulation of microcystin-RR in *Corydoras paleatus*, *Jenynsia multidentata* and *Odontesthes bonariensis*. A field and laboratory study. *Aquat. Toxicol.*, 75: 178 – 190.
- Cazenave J, Norez ML, Miceli M, Díaz MP, Wunderlin DA, Bistoni MA, 2008. Changes in the swimming activity and the glutathione – S – transferase activity of *Jenynsia multidentata* fed with microcystin-RR. *Water*

Res., 42: 1299 – 1307.

Chang TS, Jeong W, Choi SY, Yu S, Kang SW, Rhee SG, 2002. Regulation of peroxiredoxin I activity by cdc2-mediated phosphorylation. *J. Biol. Chem.*, 277: 25370 – 25376.

Chan WS, Recknagel F, Cao HQ, Park HD, 2007. Elucidation and short-term forecasting of microcystin concentrations in Lake Suwa (Japan) by means of artificial neural networks and evolutionary algorithms. *Water Res.*, 41: 2247 – 2255.

Chen L, Liu L, Huang S, 2008. Cadmium activates the mitogen – activated protein kinase (MAPK) pathway via induction of reactive oxygen species and inhibition of protein phosphatases 2A and 5. *Free Radic. Biol. Med.*, 45: 1035 – 1044.

Chen R, Chang P-A, Long D-X, Liu C-Y, Yang L, Wu Y-J, 2007. G protein β 2 subunit interacts directly with neuropathy target esterase and regulates its activity. *Int. J. Biochem. Cell Biol.*, 39: 124 – 132.

Chen T, Cui J, Liang Y, Xin X, Young DO, Chen C, Shen P, 2006. Identification of human liver mitochondrial aldehyde dehydrogenase as a potential target for microcystin-LR. *Toxicology*, 220: 71 – 80.

Chen ZH, Zhang H, Savarese TM, 1996. Gene deletion chemoselectivity: codeletion of the genes for p16 (INK4), methylthioadenosine phosphorylase, and the alpha and beta-interferons in human pancreatic cell carcinoma lines and its implications for chemotherapy. *Cancer Res.*, 56: 1083 – 1090.

Ciechanover A, Brundin P, 2003. The ubiquitin proteasome system in neurodegenerative diseases: sometimes the chicken, sometimes the egg. *Neuron*, 40(2): 427 – 446.

Clayssens S, Francois A, Chedeville A, Lavoigne A, 1995. Microcystin-LR induced an inhibition of protein synthesis in isolated rat hepatocytes. *Biochem. J.*, 306: 693 – 696.

Cohen P, 1989. The structure and regulation of protein phosphatases. *Annu. Rev. Biochem.*, 58: 453 – 508.

Collard F, Collet J, Gerin I, Veiga-da-Cunha M, Van Schaftingen E, 1999. Identification of the cDNA encoding human 6-phosphogluconolactonase, the enzyme catalyzing the second step of the pentose phosphate pathway. *FEBS Lett.*, 459: 223 – 226.

Contardo-Jara V, Pflugmacher S, Wiegand C, 2008. Multi-xenobiotic-resistance a possible explanation for the insensitivity of bivalves towards cyanobacterial toxins. *Toxicon*, 52: 936 – 943.

Craig M, Luu HA, McCready TL, Williams D, Andersen RJ, Holmes CF, 1996. Molecular mechanisms underlying the interaction of motuporin and microcystins with type – 1 and type – 2A protein phosphatases. *Biochem. Cell Biol.*, 74: 569 – 578.

Crompton M, 1999. The mitochondrial permeability transition pore and its role in cell death. *Biochem. J.*, 341: 233 – 249.

Deblois CP, Aranda-Rodriguez R, Giani A, Bird DF, 2008. Microcystin accumulation in liver and muscle of tilapia in two large Brazilian hydroelectric reservoirs. *Toxicon*, 51: 435 – 448.

Ding, WX, Shen HM, Ong CN, 2000. Critical role of reactive oxygen species and mitochondrial permeability transition in microcystin – induced rapid apoptosis in rat hepatocytes. *Hepatology*, 32: 547 – 555.

Ding WX, Shen HM, Ong CN, 2001. Pivotal role of mitochondrial Ca^{2+} in microcystin-induced mitochondrial permeability transition in rat hepatocytes. *Biochem. Biophys. Res. Commun.*, 285: 1155 – 1161.

Dittmann E, Neilan BA, Börner T, 2001. Molecular biology of peptide and polyketide biosynthesis in cyanobacteria. *Appl. Microbiol. Biotechnol.*, 57(4): 467 – 473.

Douglas P, Moorhead GB, Ye R, Lees-Miller SP, 2001. Protein phosphatases regulate DNA – dependent protein kinase activity. *J. Biol. Chem.*, 276: 18992 – 18998.

Dreumont N, Maresca A, Khandjian EW, Baklouti F, Tanguay RM, 2004. Cytoplasmic nonsense – mediated mRNA decay for a nonsense (W262X) transcript of the gene responsible for hereditary tyrosinemia, fumarylacetoacetate hydrolase. *Biochem. Biophys. Res. Commun.*, 324: 186 – 192.

Dubois T, Oudinet JP, Mira JP, Russo-Marie F, 1996. Annexins and protein kinases C. *Biochim. Biophys. Acta Mol. Cell. Res.*, 1313: 290 – 294.

Elliott WH, Elliott DC, 1997. In *Digestion and absorption of food: Biochemistry and Molecular Biology*. Oxford University Press, New York.

Faergeman NJ, Knudsen J, 1997. Role of long – chain fatty acyl-CoAs esters in the regulation of metabolism and cell signaling. *Biochem. J.*, 323: 1 – 12.

Falconer IR, Bartram J, Chorus I, Kuiper-Goodman T, Utkilen H, Burch M, Codd GA, 1999. Safe levels and safe practices. In *Toxic Cyanobacteria in Water* (I. Chorus, and J. Bartram, Eds.), pp. 155 – 178. E and FN Spon, London.

Falconer IR, 2001. Toxic cyanobacterial bloom problem in Australian waters: risks and impacts on human health. *Phycologia*, 40 (3): 228 – 233.

Falconer IR, Yeung DS, 1992. Cytoskeletal changes in hepatocytes induced by Microcystis toxins and their relation to hyperphosphorylation of cell proteins. *Chem. Biol. Interact.*, 81: 181 – 196.

Feurstein D, Holst K, Fischer A, Dietrich DR, 2009. Oatp-associated uptake and toxicity of microcystins in primary murine whole brain cells. *Toxicol. Appl. Pharmacol.*, 234: 247 – 255.

Fiaschi T, Cozzi G, Raugi G, Formigli L, Ramponi G, Chiarugi P, 2006. Redox regulation of beta-actin during integrin – mediated cell adhesion. *J. Biol. Chem.*, 281: 22983 – 22991.

Fischer WJ, Alzheimer S, Cattori V, Meier PJ, Dietrich DR, Hagenbuch B, 2005. Organic anion transporting polypeptides expressed in liver and brain mediate uptake of microcystin. *Toxicol. Appl. Pharmacol.*, 203: 257 – 263.

Fischer WJ, Dietrich DR, 2000. Pathological and biochemical characterization of microcystin – induced hepatopancreas and kidney damage in carp (*Cyprinus carpio*). *Toxicol. Appl. Pharmacol.*, 164: 73 – 81.

Fladmark KE, Brustugun OT, Mellgren G, Krakstad C, Boe R, Vintermyr OK, Schulman H, Doskeland SO, 2002. Ca²⁺/calmodulin-dependent protein kinase II is required for microcystin – induced apoptosis. *J. Biol. Chem.*, 277: 2804 – 2811.

Fontal OI, Vieytes MR, Baptista de Sousa JM, Louzao MC, Botana LM, 1999. A fluorescent microplate assay for microcystin-LR. *Anal. Biochem.*, 269: 289 – 296.

Fu WY, Yu YN, Xu LH, 2009. Identification of temporal differentially expressed protein responses to microcystin in human amniotic epithelial cells. *Chem. Res. Toxicol.*, 22(1): 41 – 51.

Gehringer MM, 2004. Microcystin-LR and okadaic acid-induced cellular effects: A dualistic response. *FEBS Lett.*, 557: 1 – 8.

Gerke V, Moss SE, 1997. Annexins and membrane dynamics. *Biochim. Biophys. Acta Mol. Cell Res.*, 1357: 129 – 154.

Gidrol X, Sabelli PA, Fern YS, Kush AK, 1996. Annexin-like protein from *Arabidopsis thaliana* rescues delta oxyR mutant of *Escherichia coli* from H₂O₂ stress. *Proc. Natl. Acad. Sci. USA*, 93: 11268 – 11273.

Gopalakrishna R, Gundimeda U, Chen ZH, 1997. Cancer preventive selenocompounds induce a specific redox modification of cysteine – rich regions in calcium-dependent isoenzymes of protein kinase C. *Arch. Biochem. Biophys.*, 348: 25 – 36.

Gorecka KM, Konopka-Postupolska D, Hennig J, Buchet R, Pikula S, 2005. Peroxidase activity of annexin 1 from *Arabidopsis thaliana*. *Biochem. Biophys. Res. Commun.*, 336: 868 – 875.

Gurbuz F, Metcalf JS, Karahan AG, Codd GA, 2009. Analysis of dissolved microcystins in surface water samples from Kovada Lake, Turkey. *Sci. Total Environ.*, 407: 4038 – 4046.

Guzman RE, Solter PF, Runnegar MT, 2003. Inhibition of nuclear protein phosphatase activity in mouse hepatocytes by the cyanobacterial toxin microcystin-LR. *Toxicon*, 41: 773 – 781.

Haarer BK, Lillie SH, Adams AEM, Magdolen V, Bandlow W, Brown SS, 1990. Purification of profilin from *Saccharomyces cerevisiae* and analysis of profiling-deficient cells. *J. Cell Biol.*, 110: 105 – 114.

Hagenbuch B, Meier PJ, 2004. Organic anion transporting polypeptides of the OATP/SLC21 family: phylogenetic classification as OATP/SLCO super family, new nomenclature and molecular/functional properties. *Pfl ü g. Arch. Eur. J. Phy.*, 447: 653 – 665.

Hanada M, Kobayashi T, Ohnishi M, Ikeda S, Wang H, Katsura K, Yanagawa Y, Hiraga A, Kanamaru R, Tamura S, 1998. Selective suppression of stress – activated protein kinase pathway by protein phosphatase 2C in mammalian cells. *FEBS Lett.*, 437: 172 – 176.

Hanada M, Ninomiya-Tsuji J, Komaki K, Ohnishi M, Katsura K, Kanamaru R, Matsumoto K, Tamura S, 2001. Regulation of the TAK1 signaling pathway by protein phosphatase 2C. *J. Biol. Chem.*, 276: 5753 – 5759.

Hayes JD, Flanagan JU, Jowsey IR, 2005. Glutathione transferases. *Annu. Rev. Pharmacol. Toxicol.*, 45: 51 – 88.

Hayward DG, Fry AM, 2006. Nek2 kinase in chromosome instability and cancer. *Cancer Lett.*, 237: 155 – 166.

Hogenboom S, Tuyp JMM, Espeel M, Koster J, Wanders RJA, Waterham HR, (2004). Human mevalonate pyrophosphate decarboxylase is localized in the cytosol. *Mol. Genet. Metab.* 81, 216 – 224.

Huang P, Xu LH, 2008. Oral administration of cyanobacterial bloom extract induced the altered expression of the PP2A, Bax and Bcl-2 in mice. *Environ. Toxicol.*, 23(6): 688 – 693.

Humble AV, Gadd GM, Codd GA, 1997. Binding of copper and zinc to three cyanobacterial microcystins quantified by different pulse polarography. *Wat. Res.*, 31(7): 1679 – 686.

Jami SK, Clark GB, Turlapati SA, Handley C, Roux SJ, Kirti PB, 2008. Ectopic expression of an annexin from *Brassica juncea* confers tolerance to abiotic and biotic stress treatments in transgenic tobacco. *Plant Physiol. Biochem.*, 46: 1019 – 1030.

Jao DL, Chen KY, 2006. Tandem affinity purification revealed the hypusine-dependent binding of eukaryotic initiation factor 5A to the translating 80S ribosomal complex. *J. Cell. Biochem.*, 97: 583 – 598.

Kaczmarek MJ, Rosenmund H, 1977. The action of human pancreatic and salivary isoamylases on starch and glycogen. *Clin. Chim. Acta*, 79: 69 – 73.

Kaplan JM, Kim SH, North KN, Rennke H, Correia LA, Tong HQ, Mathis BJ, Rodriguez-Perez JC, Allen PG, Beggs AH, Pollak MR, 2000. Mutations in ACTN4, encoding alpha-actinin-4, cause familial focal segmental glomerulosclerosis. *Nat. Genet.*, 24: 251 – 256.

Komatsu M, Furukawa T, Ikeda R, Takumi S, Nong Q, Aoyama K, Akiyama S, Keppler D, Takeuchi T, 2007. Involvement of mitogen – activated protein kinase signaling pathways in microcystin – LR – induced apoptosis after its selective uptake mediated by OATP1B1 and OATP1B3. *Toxicol. Sci.*, 97: 407 – 416.

Kondo F, Ikai Y, Oka H, Okumura M, Ishikawa N, Harada K, Matsuura K, Murata H, Suzuki M, 1992. Formation, characterization, and toxicity of the glutathione and cysteine conjugates of toxic heptapeptide microcystins. *Chem. Res. Toxicol.*, 5: 591 – 596.

Konishi T, Kato K, Araki T, Shiraki K, Takagi M, Tamaru Y, 2005. A new class of glutathione S – transferase from the hepatopancreas of the red sea bream *Pagrus major*. *Biochem. J.*, 388: 299 – 307.

Kos CH, Le TC, Sinha S, Henderson JM, Kim SH, Sugimoto H, Kalluri R, Gerszten RE, Pollak MR, 2003. Mice deficient in alpha-actinin-4 have severe glomerular disease. *J. Clin. Invest.*, 111: 1683 – 1690.

Kreplak L, Bör H, 2009. Severe myopathy mutations modify the nanomechanics of desmin intermediate filaments. *J. Mol. Bio.*, 385: 1043 – 1051.

Kujbida P, Hatanaka E, Campa A, Curi R, Farsky SHP, Pinto E, 2008. Analysis of chemokines and reactive oxygen species formation by rat and human neutrophils induced by microcystin-LA, -YR and -LR. *Toxicon*, 51: 1274 – 1280.

Kujbida P, Hatanaka E, Vinolo MA, Waismam K, Cavalcanti DM, Curi R, Farsky SH, Pinto E, 2009. Microcystins -LA, -YR, and -LR action on neutrophil migration. *Biochem. Biophys. Res. Commun.*, 382: 9 – 14.

Lankoff A, Bialczyk J, Dziga D, Carmichael WW, Lisowska H, Wojcik A, 2006. Inhibition of nucleotide excision repair (NER) by microcystin-LR in CHO-K1 cells. *Toxicon*, 48: 957 – 965.

Lankoff A, Kolataj A, 2001. Influence of microcystin-YR and nodularin on the activity of some proteolytic enzymes in mouse liver. *Toxicon*, 39: 419 – 423.

La – Salete R, Oliveira MM, Palmeira CA, Almeida J, Peixoto FP, 2008. Mitochondria a key role in microcystin-LR kidney intoxication. *J. Appl. Toxicol.*, 28: 55 – 62.

Lassing I, Schmitzberger F, Björstedt M, Holmgren A, Nordlund P, Schutt CE, Lindberg U, 2007. Molecular and structural basis for redox regulation of -actin. *J. Mol. Biol.*, 370: 331 – 348.

Lemasters JJ, Nieminen AL, Qian T, Trost LC, Elmore SP, Nishimura Y, Crowe RA, Cascio WE, Bradham CA, Brenner DA, Herman B, 1998. The mitochondrial permeability transition in cell death: A common mechanism

in necrosis, apoptosis and autophagy. *Biochim. Biophys. Acta*, 1366: 177 – 196.

Li A, Li H-Y, Jin B-F, Ye Q-N, Zhou T, Yu X-D, Pan X, Man J-H, He K, Yu M, Hu MR, Wang J, Yang SC, Shen BF, Zhang XM, 2004. A novel eIF5A complex functions as a regulator of p53 and p53-dependent apoptosis. *J. Biol. Chem.*, 279: 49251 – 49258.

Li G, Xie P, Fu J, Hao L, Xiong Q, Li H, 2008. Microcystin-induced variations in transcription of GSTs in an omnivorous freshwater fish, goldfish. *Aquat. Toxicol.*, 88: 75 – 80.

Li HH, Cai X, Shouse GP, Piluso LG, Liu X, 2007a. A specific PP2A regulatory subunit, B56[gamma], mediates DNA damage – induced dephosphorylation of p53 at Thr55. *EMBO J.*, 26: 402 – 411.

Li H, Xie P, Li G, Hao L, Xiong Q, 2009. In vivo study on the effects of microcystin extracts on the expression profiles of proto-oncogenes (c-fos, c-jun and c-myc) in liver, kidney and testis of male Wistar rats injected i.v. with toxins. *Toxicon*, 53: 169 – 175.

Li M, Satinover DL, Brautigan DL, 2007b. Phosphorylation and functions of inhibitor-2 family of proteins. *Biochemistry*, 46: 2380 – 2389.

Lindahl R, 1992. Aldehyde dehydrogenases and their role in carcinogenesis. *Crit. Rev. Biochem. Mol. Biol.*, 27: 283 – 335.

Linial M, Miller K, Scheller RH, 1989. VAT-1: an abundant membrane protein from Torpedo cholinergic synaptic vesicles. *Neuron*, 2: 1265 – 1273.

Li XY, Chung IK, Kim JI, Lee JA, 2004. Subchronic oral toxicity of microcystin in common carp (*Cyprinus carpio* L.) exposed to microcystin under laboratory conditions. *Toxicon*, 44: 821 – 827.

Li XY, Liu YD, Song LR, Liu JT, 2003. Responses of antioxidant systems in the hepatocytes of common carp (*Cyprinus carpio* L.) to the toxicity of microcystin-LR. *Toxicon*, 42: 85 – 89.

Lopez-Fanarraga M, Carranza G, Bellido J, Kortazar D, Villegas JC, Zabala JC, 2007. Tubulin cofactor B plays a role in the neuronal growth cone. *J. Neurochem.*, 100: 1680 – 1687.

Lu H, Choudhuri S, Ogura K, Csanaky IL, Lei X, Cheng X, Song PZ, Klaassen CD, 2008. Characterization of organic anion transporting polypeptide 1b2 – null mice: Essential role in hepatic uptake/toxicity of phalloidin and microcystin-LR. *Toxicol. Sci.* 103: 35 – 45.

Ma L, Li D, Wang J, He J, Yin Z, 2009. Effects of adrenergic agonists on the extrahepatic expression of vitellogenin Ao1 in heart and brain of the Chinese rare minnow (*Gobiocypris rarus*). *Aquat. Toxicol.*, 91: 19 – 25.

Malbrouck C, Trausch G, Devos P, Kestemont P, 2003. Hepatic accumulation and effects of microcystin-LR on juvenile goldfish *Carassius auratus* L. *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.*, 135: 39 – 48.

Malbrouck C, Trausch G, Devos P, Kestemont P, 2004. Effect of microcystin-LR on protein phosphatase activity and glycogen content in isolated hepatocytes of fed and fasted juvenile goldfish *Carassius auratus* L. *Toxicon*, 44: 927 – 932.

Mal é cot M, Mezhoud K, Marie A, Praseuth D, Puiseux – Dao S, Edery M, 2009. Proteomic study of the effects of microcystin-LR on organelle and membrane proteins in medaka fish liver. *Aquat. Toxicol.* 94: 153 – 161.

Marchitti SA, Deitrich RA, Vasiliou V, 2007. Neurotoxicity and metabolism of the catecholamine-derived 3,4-Dihydroxyphenylacetaldehyde and 3,4-Dihydroxyphenylglycolaldehyde: the role of aldehyde dehydrogenase. *Pharmacol. Rev.*, 59: 125 – 150.

Mattammal MB, Haring JH, Chung HD, Raghu G, Strong R, 1995. An endogenous dopaminergic neurotoxin: implication for Parkinson ' s disease. *Neurodegeneration*, 4: 271 – 281.

Maynes JT, Luu HA, Cherney MM, Andersen RJ, Williams D, Holmes CF, James MN, 2006. Crystal structures of protein phosphatase-1 bound to motuporin and dihydromicrocystin-LA: Elucidation of the mechanism of enzyme inhibition by cyanobacterial toxins. *J. Mol. Biol.*, 356: 111 – 120.

McCord JM, Fridovich I, 1988. Superoxide dismutase: the first twenty years (1968 – 1988). *Free Radic. Biol. Med.*, 5: 363 – 369.

Mezhoud K, Bauchet A.L, Château-Joubert S, Praseuth D, Marie A, Francxois JC, Fontaine JJ, Jaeg JP, Cravedi JP, Puiseux-Dao S, Edery M, 2008a. Proteomic and phosphoproteomic analysis of cellular responses in

medaka fish (*Oryzias latipes*) following oral gavage with microcystin-LR. *Toxicol.*, 51: 1431 – 1439.

Mezhoud K, Praseuth D, Puiseux-Dao S, Francxois JC, Bernard C, Edery M, 2008b. Global quantitative analysis of protein expression and phosphorylation status in the liver of the medaka fish (*Oryzias latipes*) exposed to microcystin – LR I. Balneation study. *Aquat. Toxicol.*, 86: 166 – 175.

Mikhailov A, Härmälä-Brsk é n A-S, Hellman J, Meriluoto J, Eriksson JE, 2003. Identification of ATP-synthase as a novel intracellular target for microcystin-LR. *Chem. Biol. Interact.*, 142: 223 – 237.

Monks NR, Liu S, Xu Y, Yu H, Bendelow AS, Moscow JA, 2007. Potent toxicity of the phosphatase inhibitor microcystin LR and microcystin analogues in OATPB1- and OATP1B3-expressing HeLa cells. *Mol. Cancer Ther.*, 6: 587 – 598.

Moore F, Weekes J, Hardie DG, 1991. Evidence that AMP triggers phosphorylation as well as direct allosteric activation of rat liver AMP-activated protein kinase. A sensitive mechanism to protect the cell against ATP depletion. *Eur. J. Biochem.*, 199: 691 – 697.

Moreno IM, Molina R, Jos A, Pic ó Y, Came á n AM, 2005. Determination of microcystins in fish by solvent extraction and liquid chromatography. *J. Chromatogr. A*, 1080: 199 – 203.

Moreno I, Pichardo S, Jos A, G ó mez-Amores L, Mate A, Vazquez CM, Came á n AM, 2005. Antioxidant enzymes activity and lipid peroxidation in liver and kidney of rats exposed to microcystin-LR administered intraperitoneally. *Toxicol.*, 45(4): 395 – 402.

Muhlig-Versen M, da Cruz AB, Tschape JA, Moser M, Buttner R, Athenstaedt K, Glynn P, Kretzschmar D, 2005. Loss of Swiss cheese/neuropathy target esterase activity causes disruption of phosphatidylcholine homeostasis and neuronal and glial death in adult *Drosophila*. *J. Neurosci.*, 25: 2865 – 2873.

Muslin AJ, Tanner JW, Allen PM, Shaw AS, 1996. Interaction of 14-3-3 with signaling proteins is mediated by the recognition of phosphoserine. *Cell*, 84: 889 – 897.

Muro S, P é rez B, Desviat LR, Rodr í guez-Pombo P, P é rez-Cerd á C, Clavero S, Ugarte M, 2001. Effect of PCCB gene mutations on the heteromeric and homomeric assembly of propionyl-CoA carboxylase. *Mol. Genet. Metab.*, 74: 476 – 483.

Murphy MP, 2009. How mitochondria produce reactive oxygen species. *Biochem. J.*, 417: 1 – 13.

Neves JV, Wilson JM, Rodrigues PNS, 2009. Transferrin and ferritin response to bacterial infection: The role of the liver and brain in fish. *Dev. Com. Immunol.*, 33: 848 – 857.

Nguyen A, Rothman DM, Stehn J, Imperiali B, Yaffe MB, 2004. Caged phosphopeptides reveal a temporal role for 14-3-3 in G1 arrest and S-phase checkpoint function. *Nat. Biotechnol.*, 22: 993 – 1000.

Nishizawa T, Asayama M, Fujii K, Harada K, Shriai M, 1999. Genetic analysis of the peptide synthetase genes for a cyclic heptapeptide microcystin in *Microcystis* spp.. *J. Biochem.*, 126(3): 520 – 529.

Nong Q, Komatsu M, Izumo K, Indo HP, Xu B, Aoyama K, Majima HJ, Horiuchi M, Morimoto K, Takeuchi T, 2007. Involvement of reactive oxygen species in Microcystin-LR-induced cytogenotoxicity. *Free Radic. Res.*, 41: 1326 – 1337.

O ’ Brien PJ, Siraki AG, Shangari N, 2005. Aldehyde sources, metabolism, molecular toxicity mechanisms, and possible effects on human health. *Crit. Rev. Toxicol.*, 35: 609 – 662.

Ofek P, Ben-Meir D, Kariv-Inbal Z, Oren M, Lavi S, 2003. Cell cycle regulation and p53 activation by protein phosphatase 2C alpha. *J. Biol. Chem.*, 278: 14299 – 14305.

Ohta T, Nishiwaki R, Yatsunami J, Komori A, Suganuma M, Fujiki H, 1992. Hyperphosphorylation of cytokeratins 8 and 18 by microcystin – LR, a new liver tumor promoter, in primary cultured rat hepatocytes. *Carcinogenesis*, 13: 2443 – 2447.

Oliver CJ, Shenolikar S, 1998. Physiologic importance of protein phosphatase inhibitors. *Front Biosci.*, 3: 961 – 972.

Orr PT, Jones GJ, 1998. Relationship between microcystin production and cell division rates in nitrogen – limited *Microcystis aeruginosa* cultures. *Limnol. Oceanogr.*, 43(7): 1604 – 1614.

Otsuka S, Suda S, Li R, Watanabe M, Oyaizu H, Matsumoto S, Watanabe MM, 1999. Phylogenetic relationships

between toxic and non – toxic strains of the genus *Microcystis* based on 16S to 23S internal transcribed spacer sequence. *FEMS Microbiol. Lett.*, 172(1): 15 – 21.

Oziol L, Bouaïlcha N, 2010. First evidence of estrogenic potential of the cyanobacterial hepatotoxins the nodularin-R and the microcystin-LR in cultured mammalian cells. *J. Hazard. Mater.*, 174: 610 – 615.

Paris-Palacios S, Biagianti – Risbourg S, 2006. Hepatocytes nuclear structure and subcellular distribution of copper in zebrafish *Brachydanio rerio* and roach *Rutilus rutilus* (Teleostei, Cyprinidae) exposed to copper sulphate. *Aquat. Toxicol.*, 77: 306 – 313.

Park MH, Cooper HL, Folk JE, 1981. Identification of hypusine, an unusual amino acid, in a protein from human lymphocytes and of spermidine as its biosynthetic precursor. *Proc. Natl. Acad. Sci. USA*, 78: 2869 – 2873.

Pflugmacher S, Wiegand C, Oberemm A, Beattie KA, Krause E, Codd GA, Steinberg CEW, 1998. Identification of an enzymatically formed glutathione conjugate of the cyanobacterial hepatotoxin microcystin-LR: The first step of detoxication. *Biochim. Biophys. Acta (BBA): Gen. Subj.*, 1425: 527 – 533.

Pichardo S, Jos A, Zurita JL, Salguero M, Camean AM, Repetto G, 2007. Acute and subacute toxic effects produced by microcystin-YR on the fish cell lines RTG-2 and PLHC-1. *Toxicol. Vitro*, 21: 1460 – 1467.

Pouria S, de Andrade A, Barbosa J, Cavalcanti RL, Barreto VT, Ward CJ, Preiser W, Poon GK, Neild GH, Codd GA, 1998. Fatal microcystin intoxication in haemodialysis unit in Caruaru, Brazil. *The Lancet*, 352(9121): 21 – 26.

Ratner N, Mahler HR, 1983. Structural organization of filamentous proteins in postsynaptic density. *Biochemistry*, 22: 2446 – 2453.

Rhee HJ, Kim GY, Huh JW, Kim SW, Na DS, 2000. Annexin I is a stress protein induced by heat, oxidative stress and a sulfhydryl-reactive agent. *Eur. J. Biochem.*, 267: 3220 – 3225.

Robinson WG, Coon MJ, 1957. Purification and properties of beta-hydroxyisobutyric dehydrogenase. *J. Biol. Chem.*, 225: 511 – 521.

Sacre SM, Moss SE, 2002. Intracellular localization of endothelial cell annexins is differentially regulated by oxidative stress. *Exp. Cell Res.*, 274: 254 – 263.

Sicinska P, Bukowska B, Michalowicz J, Duda W, 2006. Damage of cell membrane and antioxidative system in human erythrocytes incubated with microcystin-LR in vitro. *Toxicon*, 47: 387 – 397.

Singh DP, Tyagi MB, Kumar A, Thakur JK, Kumar A, 2001. Antialgal activity of a hepatotoxin-producing cyanobacterium, *Microcystis aeruginosa*. *World J. Microbiol. Biotech.*, 17(1): 15 – 22.

Song LR, Chen W, Peng L, Wan N, Gan NQ, Zhang XM, 2007. Distribution and bioaccumulation of microcystins in water columns: A systematic investigation into the environmental fate and the risks associated with microcystins in Meiliang Bay, Lake Taihu. *Water Res.*, 41: 2853 – 2864.

Soares RM, Yuan M, Servaites JC, Delgado A, Magalhaes VF, Hilborn ED, Carmichael WW, Azevedo SM, 2006. Sublethal exposure from microcystins to renal insufficiency patients in Rio de Janeiro, Brazil. *Environ. Toxicol.*, 21: 96 – 103.

Sobue K, Kanda K, 1989. Alpha – actinins, caldesmon (brain spectrin or fodrin), and actin participate in adhesion and movement of growth cones. *Neuron*, 3: 311 – 319.

Taylor CA, Senchyna M, Flanagan J, Joyce EM, Clich é DO, Boone AN, Culp-Stewart S, Thompson JE, 2004. Role of eIF5A in TNF-alpha-mediated apoptosis of lamina cribrosa cells. *Invest. Ophthalmol. Visual Sci.*, 45: 3568 – 3576.

Talpade DJ, Greene JG, Higgins Jr. DS, Greenamyre JT, 2000. In vivo labeling of mitochondrial complex I (NADH:ubiquinone oxidoreductase) in rat brain using [(3)H]dihydrorotenone. *J. Neurochem.*, 75: 2611 – 2621.

Tanaka T, Akatsuka S, Ozeki M, Shirase T, Hiai H, Toyokuni S, 2004. Redox regulation of annexin 2 and its implications for oxidative stress – induced renal carcinogenesis and metastasis. *Oncogene*, 23: 3980 – 3989.

Tildon JT, Sevdalian DA, 1972. CoA transferase in the brain and other mammalian tissues. *Arch. Biochem. Biophys.*, 148: 382 – 390.

- Tillett D, Dittmann E, Erhard M, von Dohren H, Borner T, Neilan BA, 2000. Structural organization of microcystin biosynthesis in *Microcystis aeruginosa* PCC7806: an integrated peptide – polyketide synthetase system. *Chem. Biol.*, 7(10): 753 – 764.
- Toh BH, Gallichio HA, Jeffrey PL, Livett BG, Muller HK, Cauchi MN, Clarke FM, 1976. Anti – actin stains synapses. *Nature*, 264: 648 – 650.
- Tomsig JL, Sohma H, Creutz CE, 2004. Calcium-dependent regulation of tumour necrosis factor – alpha receptor signalling by copine. *Biochem. J.*, 378: 1089 – 1094.
- Toomey BH, Epel D, 1995. A multixenobiotic transporter in *Urechis caupo* embryos: Protection from pesticides? *Mar. Environ. Res.*, 39: 299 – 302.
- Tsujimoto Y, Shimizu S, 2002. The voltage-dependent anion channel: an essential player in apoptosis. *Biochimie*, 84: 187 – 193.
- Ueno Y, Nagata S, Tsutsumi T, Hasegawa A, Watanabe MF, Park HD, Chen G, Yu SZ, 1996. Detection of microcystin, a blue-green algal hepatotoxin, in drinking water sampled in Haimen and Fusui, endemic areas of primary liver cancer in China, by highly sensitive immunoassay. *Carcinogenesis*, 17(6): 1317 – 1321.
- Wang DZ, Lin L, Chan LL, Hong HS, 2009. Comparative studies of four protein preparation methods for proteomic study of the dinoflagellate *Alexandrium* sp. using two-dimensional electrophoresis. *Harmful Algae* 8: 685 – 691.
- Wang L, Liang XF, Liao WQ, Lei LM, Han BP, 2006. Structural and functional characterization of microcystin detoxification-related liver genes in a phytoplanktivorous fish, Nile tilapia (*Oreochromis niloticus*). *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.*, 144: 216 – 227.
- Wang M, Chan LL, Si M, Hong H, Wang D, 2010. Proteomic analysis of hepatic tissue of zebrafish (*Danio rerio*) experimentally exposed to chronic microcystin-LR. *Toxicol. Sci.*, 113: 60 – 69.
- Wang W, Ding J, Allen E, Zhu P, Zhang L, Vogel H, Yang Y, 2005. Gigaxonin interacts with tubulin folding cofactor B and controls its degradation through the ubiquitin – proteasome pathway. *Curr. Biol.*, 15: 2050 – 2055.
- Wei Y, Weng D, Li F, Zou X, Young DO, Ji J, Shen P, 2008. Involvement of JNK regulation in oxidative stress – mediated murine liver injury by microcystin-LR. *Apoptosis*, 13: 1031 – 1042.
- Weng D, Lu Y, Wei Y, Liu Y, Shen P, 2007. The role of ROS in microcystin-LR-induced hepatocyte apoptosis and liver injury in mice. *Toxicology*, 232: 15 – 23.
- Wickstrom ML, Khan SA, Haschek WM, Wyman JF, Eriksson JE, Schaeffer DJ, Beasley VR, 1995. Alterations in microtubules, intermediate filaments, and microfilaments induced by microcystin – LR in cultured cells. *Toxicol. Pathol.*, 23: 326 – 337.
- Wierzbicki AS, Lloyd MD, Schofield CJ, Feher MD, Gibberd FB, 2002. Refsum ' s disease: a peroxisomal disorder affecting phytanic acid alpha-oxidation. *J. Neurochem.*, 80: 727 – 735.
- Xing ML, Wang XF, Xu LH, 2008. Alteration of proteins expression in apoptotic FL cells induced by MCLR. *Environ. Toxicol.*, 23: 451 – 458.
- Xing YN, Li Z, Chen Y, Stock JB, Jeffrey PD, Shi YG, 2008. Structural mechanism of demethylation and inactivation of protein phosphatase 2A. *Cell*, 133: 154 – 163.
- Xing, Y, Xu Y, Chen Y, Jeffrey PD, Chao Y, Lin Z, Li Z, Strack S, Stock JB, Shi Y, 2006. Structure of protein phosphatase 2A core enzyme bound to tumor-inducing toxins. *Cell*, 127: 341 – 353.
- Yeung K, Seitz T, Li S, Janosch P, McFerran B, Kaiser C, Fee F, Katsanakis KD, Rose DW, Mischak H, Sedivy JM, Kolch W, 1999. Suppression of Raf-1 kinase activity and MAP kinase signalling by RKIP. *Nature*, 401: 173 – 177.
- Zegura B, Zajc I, Lah TT, Filipic M, 2008. Patterns of microcystin-LR induced alteration of the expression of genes involved in response to DNA damage and apoptosis. *Toxicon*, 51: 615 – 623.

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库